

## Research Supporting the Importance of DHA for Adult Health

### Brain and Eye Health

1. Salem, Jr. N, et al. Mechanisms of action of docosahexaenoic acid in the nervous system. *Lipids*, 2001. 36:945-59.
2. Marszalek JR, et al. Docosahexaenoic Acid, Fatty-Acid-Interacting Proteins, and Neuronal Function: Breastmilk and Fish are Good for You. *Annu Rev Cell Dev Bio*, 2005. 21:633-57.
3. Heude B, et al. Cognitive decline and fatty acid composition of erythrocyte membranes - The EVA Study. *Am J Clin Nutr*, 2003. 77:803-808.
4. Soderberg, M., et al. Fatty acid composition of brain phospholipids in aging and in Alzheimer's disease. *Lipids*. 1991;26:421-425.
5. Yazawa, K. "Clinical experience with docosahexaenoic acid in demented patients," International Conference on Highly Unsaturated Fatty Acids in Nutrition and Disease Prevention, Barcelona, Spain.1996.
6. Kyle, D.J., et al. Low serum docosahexaenoic acid is a significant risk factor for Alzheimer's dementia. *Lipids*. 1999;34:S245.
7. Morris MC, et al. Consumption of fish and n-3 fatty acids and risk of incident Alzheimer disease. *Arch Neurol*, 2003. 60:940-6.
8. Maclean CH, et al. Effects of omega-3 fatty acids on cognitive function with aging, dementia, and neurological diseases. *Evid Rep Technol Assess (Summ)*, 2005. 1-3.
9. Hashimoto M, et al. Chronic administration of docosahexaenoic acid ameliorates the impairment of spatial cognition learning ability in amyloid beta-infused rats. *J Nutr*, 2005. 135:549-555
10. Lukiw WJ, et al. A role for docosahexaenoic acid-derived neuroprotectin D1 in neural cell survival and Alzheimer disease. *J of Clin Investigation*, 2005. 115:2774-2783.
11. Lim GP, et al. A diet enriched with the omega-3 fatty acid docosahexaenoic acid reduces amyloid burden in an aged Alzheimer mouse model. *J Neurosci*, 2005. 25(13).
12. Florent S, et al. Docosahexaenoic acid prevents neuronal apoptosis induced by soluble amyloid-B oligomers. *J Neurochem*, 2006. 96(2):385-95.
13. SanGiovanni JP, et al. The role of omega-3 long-chain polyunsaturated fatty acids in health and disease of the retina. *Prog Retin Eye Res*, 2005. 24:87-138.
14. Hoffman, D.R. and D.G. Birch. Docosahexaenoic acid in red blood cells of patients with x-linked retinitis pigmentosa. *Invest Ophthalmol Vis Sci*. 1995. 36:1009-1018.
15. Schaefer, E.J., et al. Red blood cell membrane phosphatidylethanolamine fatty acid content in various forms of retinitis pigmentosa. *J Lipid Res*. 1995. 36:1427-1433.
16. Harding, C.O., et al. Docosahexaenoic acid and retinal function in children with long-chain 3-hydroxyacyl-CoA dehydrogenase deficiency. *J Inherit Metab Dis*. 1999. 22(3): p. 276-80.

### Research Supporting the Cardioprotective Benefits of DHA

1. O'Keefe, J.H. and W.S. Harris. From Inuit to implementation: omega-3 fatty acids come of age. *Mayo Clin Proc*, 2000. 75:607-614.
2. Valagussa, F.E.A., et al. Dietary supplementation with n-3 polyunsaturated fatty acids and vitamin E after myocardial infarction: results of the GISSI-Prevenzione trial. Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto miocardico. *Lancet*, 1999. 354:447-445.
3. Albert, C.M., et al. Fish consumption and risk of sudden cardiac death. *JAMA*, 1998. 279:23-28.
4. Daviglius, M.L., et al. Fish consumption and the 30-year risk of fatal myocardial infarction. *N Engl J Med*, 1997. 336:1045-1053.
5. Siscovick, D.S., et al. Dietary intake and cell membrane levels of long-chain n-3 polyunsaturated fatty acids and the risk of primary cardiac arrest. *JAMA*, 1995. 274:1363-1367.
6. Singh, R.B., et al. Randomized, double-blind, placebo-controlled trial of fish oil and mustard oil in patients with suspected acute myocardial infarction: the Indian experiment of infarct survival – 4. *Cardiovasc Drugs Ther*, 1997. 11:485-491.
7. Dolecek, T.A. Epidemiological evidence of relationships between dietary polyunsaturated fatty acids and mortality in the Multiple Risk Factor Intervention Trial. *Proc Soc Exp Biol Med*, 1992. 200:177-182.
8. Kromhout, D., et al. The inverse relation between fish consumption and 20-year mortality from coronary heart disease. *N Engl J Med*, 1985. 312:1205-1209.
9. Benisek, D. and D. Kyle. Socioeconomic differences in the consumption of polyunsaturated fatty acids in the United States. *J Am Coll Nutr*, 1999. 18(5):543-544.
10. American Heart Association Home Page. [www.americanheart.org](http://www.americanheart.org). 2000.
11. Agren, J.J., et al. Fish diet, fish oil and docosahexaenoic acid rich oil lower fasting and postprandial plasma lipid levels. *Eur J Clin Nutr*, 1996. 50:765-771.
12. Nelson, G.J., et al. The effect of dietary docosahexaenoic acid on plasma lipoprotein and tissue fatty acid composition in humans. *Lipids*, 1997. 32(11):1137-46.
13. Conquer, J.A. and B.J. Holub. Supplementation with an algae source of docosahexaenoic acid increases (n-3) fatty acid status and alters selected risk factors for heart disease in vegetarian subjects. *J Nutr*, 1996. 126:3032-3039.
14. Innis, S.M. and J.N. Hansen. Plasma fatty acid responses, metabolic effects, and safety of microalgal and fungal oils rich in arachidonic and docosahexaenoic acid in healthy adults. *Am J Clin Nutr*, 1996. 64:159-67.
15. Davidson, M.H., et al. Effects of docosahexaenoic acid on serum lipoproteins in patients with combined hyperlipidemia: a randomized, double-blind, placebo-controlled trial. *J Am Coll Nutr*, 1997. 16(3):236-243.
16. Mori, T.A., et al. Purified eicosapentaenoic and docosahexaenoic acids have different effects on serum lipids and lipoproteins, LDL particle size, glucose, and insulin in mildly hyperlipidemic men. *Am J Clin Nutr*, 2000. 71:1085-94.

17. Mori, T.A., et al. Docosahexaenoic acid but not eicosapentaenoic acid lowers ambulatory blood pressure and heart rate in humans. *Hypertension*, 1999. 34:253-260.